Josera.
we care, you grow

Combivit

Supplementary feed for breeding animals for temporary, additional application due to high performance requirements

Special situations require special solutions. High performance combined with stressful conditions — such as regrouping, drying-off, calving, transport, or high infection pressure — are such situations. To prevent performance drops in these cases, it is beneficial to give the metabolism a "positive shock." A short-term vitamin boost briefly disrupts homeostasis and helps the metabolism cope with extreme situations.

Combivit contains a high-quality vitamin profile, including B vitamins that support metabolic function.
The result is strong resilience during times of stress and high demand.

As a result, Combivit leads to:
- Reduced performance losses in extreme situations
- Increased safety
- A strong immune system and metabolism

Main applications

- early dry period
- preparation for birth
- lactation
- young cattle
- fattening cattle
- suckling cows

Product advantages

- ✓ high palatability
- ✓ strong immune system
- ✓ stable metabolism



PERFORMANCE

Combivit

Supplementary feed for breeding animals for temporary, additional application due to high performance requirements

Product features:
- high value of vitamins
- B-vitamins for the relief of the metabolism

Application recommendation:
Concentrated supplement for temporary, additional application due to high performance, stress, high risk of infection and other extraordinary exposures.

Feeding recommendation:

a) Dairy cows
- For drying up, 200g per day for 3 days
- 3 weeks before calving, 200g per day for 3 days
- 3 days before calving to 7 days after, 200g per day
- At the first heat, 200g per day for 3 days
- Before breeding, 200g per day for 3 days

b) Breeding bulls
- Monthly 200g per day for 3 days

c) Calves for weaning
- 50g per day for 3 - 5 days

d) Sheep during breeding season and nursing
- 50g per day for 3 - 5 days

e) In problem situations such as birth, feed change, barn change, transportation, high infection risks, etc.
- 200g per day for 3 - 5 days
- Cattle 100g each
- Breeding pigs 50g each
- Horses, breeding mares 100g each
- Calves, ewes 50g each

Instructions for proper use:
This complementary feed has a higher content of vitamins than complete feed. Therefore it must be given to dairy cows, heifers, cattle and horses only in a quantity of 1% per daily ration.

Composition:
wheat | wheat bran | apple pomace | yeast | sodium chloride | calcium carbonate

Analytical constituents:

14,00 %	crude protein	3,00 %	crude fat
7,50 %	crude fibre	20,00 %	crude ash
0,80 %	lysine	2,00 %	calcium
0,50 %	phosphorus	2,00 %	sodium

Nutritional additives per 1 kg:

10.000,0 mg	vitamin E/all-rac-alpha-tocopheryl acetate (3a700)
5.000,0 mg	vitamin C (3a300)
2.350,0 mg	vitamin B1 (3a821)
1.125,0 mg	vitamin B2 (3a825ii)
1.000,0 mg	vitamin B6/ pyridoxine hydrochloride (3a831)
15.000,0 mcg	vitamin B12 (3a835 / 3a836 / 3a837)
6.750,0 mg	niacinamide (3a315)
3.750,0 mg	calcium D-pantothenate (3a841)
150,0 mg	folic acid (3a316)
5.000,0 mg	choline chloride (3a890)
100.000,0 mcg	biotin (3a880)

Note: The best before date requires hygienic, cool, and dry storage. Josera does not take responsibility for incorrect storage nor improper use.

20 kg

PERFORMANCE

Please use the JOSERA advisory service for targeted advice.

Josera GmbH | Industriegebiet Süd | 63924 Kleinheubach | Germany
Tel.: +49 (0) 9371 940-0 | Fax: +49 (0) 9371 940-149 | export@josera.de | www.josera-agriculture.com | α DE-BY-1-00035

EN044

Special product